

Title	Prelithification and synlithification tectonic foliation development in a clastic sedimentary sequence
Authors	Meere, Patrick A.;Mulchrone, Kieran F.;McCarthy, David J.;Timmerman, Martin J.;Dewey, John F.
Publication date	2016-04
Original Citation	Meere, P. A., Mulchrone, K. F., McCarthy, D. J., Timmerman, M. J. and Dewey, J. F. (2016) 'Prelithification and synlithification tectonic foliation development in a clastic sedimentary sequence', [2016] <i>Geology</i> , 44(4), pp. 291-294. doi:10.1130/G37587.1
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://pubs.geoscienceworld.org/gsa/geology/article/44/4/291/132050/prelithification-and-synlithification-tectonic - 10.1130/G37587.1
Rights	© 2016 Geological Society of America.
Download date	2023-05-05 04:23:18
Item downloaded from	http://hdl.handle.net/10468/6240



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

1 Pre/syn-lithification tectonic foliation development in a
2 clastic sedimentary sequence.

3 **Patrick A. Meere^{1*}, Kieran F. Mulchrone², Dave J. McCarthy³, Martin**
4 **Timmerman⁴, and John F. Dewey⁵**

5 *¹School of Biological, Earth & Environmental Science, Environmental Research Institute,*
6 *University College Cork, Cork T12 YN60, Ireland*

7 *²Department of Applied Mathematics, School of Mathematics, University College Cork,*
8 *Cork T12 YN60, Ireland*

9 *³British Geological Survey, Murchison House, W Mains Rd, Edinburgh EH9 3LA, UK*

10 *⁴Institut für Erd-und Umweltwissenschaften, Universität Potsdam, Karl-Liebknecht-*
11 *Strasse 24-25, 14476 Potsdam-Golm, Germany*

12 *⁵University College, High Street, Oxford OX1 4BH, UK*

13 *E-mail: p.meere@ucc.ie

14 **ABSTRACT**

15 The current view regarding the timing of regionally developed penetrative
16 tectonic fabrics in sedimentary rocks is that their development postdates lithification of
17 those rocks. In this case, fabric development is achieved by a number of deformation
18 mechanisms including grain rigid body rotation, crystal-plastic deformation and pressure
19 solution. The latter is believed to be the primary mechanism responsible for the domainal
20 structure of cleavage in low-grade metamorphic rocks. In this study we combine field
21 observations with strain studies to characterize considerable (>50%) Acadian crustal
22 shortening in a Devonian clastic sedimentary sequence from southwest Ireland. Despite

these high levels of shortening there is a marked absence of the domainal cleavage structure and intra-clast deformation, which are expected with this level of deformation. Fabrics in these rocks are predominantly a product of rigid body rotation and repacking of extra-formational clasts during deformation of a clastic sedimentary sequence before lithification was complete.

INTRODUCTION

Attempting to understand the key physical/chemical processes of tectonic foliation formation has occupied the minds of some of the leading geologists for nearly 200 years (Darwin, 1846; Sorby, 1849) with answers to some fundamental questions still outstanding. Research since the early seventies has emphasized the central role pressure dissolution plays in the formation of tectonic cleavage (Wood, 1974, Vernon, 1998). As a consequence, cleavage foliations are typically domainal with alternating phyllosilicate-rich dissolution cleavage domains and lithon domains of relatively un-deformed host lithology (Powell, 1979; Borradaile et al., 1982; Vernon, 1998). Deformation mechanisms involved in the formation of these fabrics include grain rigid body rotation producing grain shape preferred orientation (GSPO), crystal-plastic deformation and pressure dissolution (Vernon, 1998). The current orthodoxy is that these processes predominantly operate to produce a slaty cleavage after the host lithology has become fully lithified (Vernon, 2004, *and references therein*). While there have been advocates for pre-lithification development of tectonic fabrics (Maxwell, 1962; Alterman, 1973) these examples are viewed as 'local' aberrations that are not regionally significant (Geiser, 1975). However in recent years there has been a growing awareness of the role of 'lateral compaction' in producing a distributed shortening strain in partially lithified

sediments (Paterson and Tobisch, 1993; Henry et al., 2003; Butler and Paton, 2010, Alsop and Marco, 2014). Butler and Paton (2010) estimated up to 25% distributed longitudinal strain in a gravity driven thrust system from the Orange Basin offshore Namibia. Here we describe a Devonian clastic sedimentary sequence from southern Ireland that has experienced considerable shortening associated with tectonic foliation development yet exhibits minimal evidence of structures typically associated with deformation of lithified rocks. Evidence is presented that regional tectonic shortening was achieved by translation and rigid body rotation of clasts with possible concomitant sediment dewatering of a not fully lithified sedimentary sequence.

BACKGROUND GEOLOGY

The Dingle Peninsula of southwest Ireland consists of a series of distinct tectono-stratigraphic units representing alternating periods of localized crustal extension and compression extending from the late Silurian to the early Carboniferous. One of these, the Dingle Group represents the early continental infilling of the Lower Devonian Dingle Basin. This basin extends for ~60 km along the axis of the Dingle Peninsula and has been described as a pull-apart structure within the Caledonian Iapetus Suture Zone (Todd, 2000). The basin fill, the Dingle Group, is predominantly fluvial and includes two marginal conglomerate units, the Glashabeg Formation preserved along the northern margin of the basin and Trabeg Formation along the southern margin (Horne, 1974). This study focuses on the Glashabeg Formation in the Wine Strand area (52.17871°N, 10.38488°W) on the northwestern side of the peninsula (Fig. 1). Compositionally the Glashabeg Formation consists of a series of fining-upward cycles consisting of polymict basal conglomerates overlain by red sandstones, siltstones and mudstones. The

conglomerates predominantly consist of volcanic and siltstone extra-formational clasts with variable amounts of jasper, vein quartz and critically intra-formational 'rip up' mud/siltstone clasts set in a very coarse grained sandstone matrix (Figs. 2a–2d). After deposition, this basin fill was deformed by the mid-Devonian Acadian orogenic event (Meere and Mulchrone, 2006) leading to regional fabric development, folding and localized reverse faulting. The study area sits close to the core of an open and upright Acadian syncline, the Ballyferriter Syncline, which plunges gently to the northeast. A penetrative tectonic fabric (Fig. 2e) transects the syncline axis by $\sim 14^\circ$ anticlockwise (Fig. 1) consistent with regional dextral Acadian transpression (Meere and Mulchrone, 2006). The xy (flattening) principle planes of finite strain (R_s) derived from oblate reduction spots lie parallel to the cleavage fabric with a mean xz R_s value of 2.73 ± 0.25 (Meere and Mulchrone, 2006). This equates to $\sim 50\%$ bulk shortening, assuming constant volume deformation, or $\sim 65\%$ shortening, assuming a volume loss deformation process. The maximum principle strain x axis of the xy section ellipses consistently pitch steeply in the cleavage plane indicating a component of sub-vertical thickening associated with tectonic shortening. The deformation occurred under very low grade (sub-greenschist) metamorphic conditions with no evidence of metamorphic mineral growth. Palynomorphs taken from Dingle Group rocks are black in color (Higgs et al., 2014) indicating a thermal alteration index (TAI) of 4.5–5 indicative of maximum paleo-temperatures in excess of 250°C but below greenschist metamorphic facies conditions.

FIELD EVIDENCE

A number of features have been recognized in Glashabeg Formation lithologies that are unusual for rocks that have undergone such significant levels of tectonic shortening;

(1) With the exception of some very localized Mode 1 fracturing, there is an absence of intra clast deformation in conglomerate extra-formational clasts (Figs. 1a and 1b). There is no evidence of pressure dissolution indenting at clast/clast contact points. Isolated extra-formational clasts in matrix-rich conglomerates display strong 'wrap around' fabrics developed in the vicinity of the clast indicating more competent behavior with respect to the enclosing matrix during deformation (Figs. 2a and 2b). In addition, there is no evidence of such features as 'rolling structures' (Van den Driessche and Brun, 1987) indicating clast rotation that would be expected with ductile deformation of a fully lithified conglomerate. Similar fabrics have been described in the Lafonia Diamictite of the Falkland Islands (Curtis and Hyam, 1998).

(2) In sharp contrast, intra-formational mud and fine siltstone 'rip up' clasts have behaved less competently during deformation with clast/matrix boundaries often displaying convex inward 'bulging' structures (Fig. 2c) (Waldron and Gagnon, 2011). While this indicates that the 'rip-up' clasts were less competent than the surrounding matrix, it also requires that both materials were in a less competent weakly lithified state during deformation. Where competent extra-formational clasts are in direct contact with 'rip-up' clasts they are seen to project into the less competent mudstone/siltstone of the 'rip-up' clasts (Fig. 2b). Intra-formational clasts also consistently show very strong alignment parallel to the tectonic fabric, even in areas where there is significant discordance between this fabric and the primary bedding fabric (Fig. 2d).

(3) Overall, finer grained siltstone and mudstone lithologies exhibit a high level of less competent behavior during deformation. High amplitude mullion structures are typically developed at mudstone/conglomerate contacts (Figs. 2e and 2f) with the less competent mudstone cusps projecting into the more competent conglomerates. This mullion lineation is parallel to the regional bedding/cleavage intersection lineation.

(4) On a microscopic scale there is a marked absence of a pervasive domainal microstructure, grain flattening and pressure solution seam development. The absence of these microstructure is indicative of soft-sediment deformation fabrics the development of which is characterized by rigid body grain rotation (Waldron and Gagnon, 2011; Alsop and Marco, 2014). Qualitative element concentration maps of the finer grained lithologies were made using a JEOL JXA-8200 electron probe micro-analyzer at the Universität Potsdam (Germany) which is equipped with five wavelength-dispersive spectrometers and operated at 15 kV accelerating voltage and 35 nA sample current. Critically, these maps confirm the absence of Si depleted seams as described by Meere et al. (2013). Structures indicative of intra-crystalline deformation such as pervasive undulose extinction, sub-grain development or any recrystallization mechanisms are absent and ought to be present in the case of pervasive deformation of lithified sedimentary rocks. Where dissolution seam development occurs it is very localized, typically developing in intra-formational mudstone clasts, due to high mean stress concentrations at the apices of extra-formational clasts projecting into the less competent 'rip-up' clast material (Fig. 3b). Boundaries between siltstones and coarse sandstones are often characterized by isolated sandstone clasts completely embedded in siltstone (Fig. 3a).

136 STRAIN ANALYSIS

137 Finite strain (R_s) estimates obtained from reduction spots were compared to those
138 derived in this study from siltstone, sandstone and conglomerate samples using the R_f/ϕ
139 mean radial length (MRL) (Mulchrone et al., 2003) strain analysis method. This method
140 assumes passive clast/matrix material behavior as well as an initial random distribution of
141 clast orientations and a radial symmetry of clast axial ratios. With increasing departure
142 from these assumptions the MRL method will increasingly underestimate the true R_s
143 value. A minimum of 150 clast aspect ratios/orientations were collected from each
144 analyzed sample to reduce error associated with the finite strain estimates (Meere and
145 Mulchrone, 2003). Data were collected from shallow dipping units where the tectonic
146 fabric was $\sim 90^\circ$ to bedding and where there was good control on finite strain from high
147 quality reduction spot data in adjacent mudstones and siltstones. Data has been extracted
148 using semi-automatic analysis of digital images (Mulchrone et al., 2013). Previous
149 studies on the reduction spots show marked discontinuities in the curvature of the
150 reduction spot boundaries between fine-grained and coarse-grained siltstone components.
151 This indicates differential shortening within these lithologies during cleavage
152 development which in turn indicates they developed before deformation and are as such
153 valid finite strain markers (Meere et al. 2008).

154 Results for all sediment grain sizes (Fig. 4a) clearly show significant
155 underestimates of finite strain with respect to the reduction spot data ($R_s = 2.73 \pm 0.25$)
156 strongly indicating that the assumptions of MRL, principally passive clast/matrix
157 behavior are not valid. In all cases the finite strain x axis is closely aligned to the trace of
158 the cleavage fabric (S_1). By contrast, the intra-formational 'rip-up' clast sample gives the

highest MRL strain estimate ($R_s = 2.2$). Field evidence which suggests less competent behavior is consistent with finite strain estimates that more closely approximate the true strain value.

STRAIN MODELING

Structures observed in the field strongly indicate that conglomerates reacted to deformation in the unconsolidated state. Therefore associated clast fabrics cannot be explained in terms of traditional passive behavior (Mulchrone et al., 2003). In the unconsolidated state clasts behave like rigid inclusions by comparison with the enclosing matrix. The motion of rigid inclusions with no-slip at the boundary is well understood (Jeffery, 1922) and it is possible to relate distributions of clast long axis orientations to finite strain and strain history (Mulchrone, 2007a). Models of the case of rigid inclusions with slip on the boundary have also been developed (Mulchrone, 2007b). By deriving probability distribution functions for both no-slip and slip boundary conditions, maximum likelihood methods allow for estimation and comparison of finite strain from long axis distributions (Mulchrone and Meere, 2015) for both cases. Therefore an appropriate model of clast behavior can be determined by calculating clast fabric intensity under these two different boundary conditions and comparing the results with natural data.

The axial ratios and orientations of 315 conglomerate clasts from the Glashabeg Formation were measured in a section normal to bedding and the tectonic fabric. The data were analyzed assuming pure shear, and both 'rigid no-slip' and 'rigid slip' boundary conditions. The results are summarized as a plot of fabric intensity versus bulk strain (R_s) (Fig. 4b). Under the assumption of 'rigid no-slip' it takes a finite strain of $R_s > 14.0$ to

produce the observed clast fabric intensity whereas assuming 'rigid slip' behavior the observed distribution is explained by a finite strain of $R_s = 2.4$ which is close to the bulk strain estimate derived from reduction spots.

CONCLUSIONS

A number of lines of evidence from the Glashabeg Formation support the contention that these rocks were deformed before the process lithification was complete. These include;

- (1) An absence of a pervasive dissolution seam (Si depleted) fabrics.
- (2) A spectrum of clast/matrix interactions from rigid extra-formational clast behaviors (e.g., fabric wrapping around clasts) to less competent behaviors (e.g., bulging) for less competent intra-formational clasts.
- (3) An absence of 'rolling structures' indicating clast rotation in a lithified matrix during deformation.
- (4) The presence of high amplitude lobate mullion structures are developed at mudstone/conglomerate contacts
- (5) Strain analysis results for extra-formational clasts clearly show significant underestimates of finite strain while results for the more incompetent 'rip up' clasts yield higher estimates ($R_s = 2.2$) closer to the true strain values from reduction spot data ($R_s = 2.73 \pm 0.25$).
- (6) Strain modeling indicates that the observed clast fabric intensities are consistent with 'rigid slip' behavior of extra-formational clasts in a weak matrix.

The deformation of poorly lithified sediments proposed in this study is consistent with the close temporal proximity of the deposition of the Lower Devonian Dingle Group

sediments in the Dingle Basin and their subsequent deformation by the mid-Devonian
Acadian event in southwest Ireland. This study revives the argument for a mechanism of
developing a well-defined tectonic fabric prior to lithification (Maxwell, 1962) and
requires geologists to consider the possibility of such a mechanism contributing to
tectonic strain in a range of geological settings. It also has implications for sediment
mobility during deformation. This includes the preferential exploitation of pre-existing
tectonic fabrics by emplacement of clastic dikes (Dewey and Ryan, 1990, Phillips and
Alsop, 2000). These results also highlight the importance of demonstrating passive
clast/matrix behavior when deriving meaningful finite strain estimates using most
conventional strain analysis techniques based on clast population behavior during
deformation.

REFERENCES CITED

- Alsop, G.I., and Marco, S., 2014, Fold and fabric relationships in temporally and spatially
evolving slump systems: A multi-cell flow model: *Journal of Structural Geology*,
v. 63, p. 27–49, doi:10.1016/j.jsg.2014.02.007.
- Alterman, I., 1973, Rotation and dewatering during slaty cleavage formation: some new
evidence and interpretations: *Geology*, v. 1, p. 33–36, doi:10.1130/0091-
7613(1973)1<33:RADDSC>2.0.CO;2.
- Borradaile, G.J., Bayly, M.B., and Powell, C.M.A., eds., 1982, *Atlas of deformational
and metamorphic rock fabrics*: Heidelberg, New York, Springer, doi:10.1007/978-3-
642-68432-6.

- 226 Butler, R.W.H., and Paton, D.A., 2010, Evaluating lateral compaction in deepwater fold
227 and thrust belts: How much are we missing from “nature’s sandbox”? GSA Today,
228 v. 20, p. 4–10, doi:10.1130/GSATG77A.1.
- 229 Curtis, M.L., and Hyam, D.M., 1998, Late Palaeozoic to Mesozoic structural evolution of
230 the Falkland Islands: A displaced segment of the Cape Fold Belt: Journal of the
231 Geological Society, v. 155, p. 115–129, doi:10.1144/gsjgs.155.1.0115.
- 232 Darwin, C., 1846, Geological Observations on South America, Being the Third Part of
233 the Geology of the Voyage of the “Beagle” During the Years 1832 to 1836: London,
234 Smith-Elder.
- 235 Dewey, J.F., and Ryan, P.D., 1990, The Ordovician evolution of the South Mayo Trough,
236 western Ireland: Tectonics, v. 9, p. 887–901, doi:10.1029/TC009i004p00887.
- 237 Geiser, P.A., 1975, Slaty cleavage and the dewatering hypothesis — An examination of
238 some critical evidence: Geology, v. 3, p. 717–720, doi:10.1130/0091-
239 7613(1975)3<717:SCATDH>2.0.CO;2.
- 240 Henry, P., Jouniaux, L., Screaton, E.J., Hunze, S., and Saffer, D.M., 2003, Anisotropy of
241 electrical conductivity record of initial strain at the toe of the Nankai accretionary
242 prism: Journal of Geophysical Research, v. 108, B9, p. 2407–2418,
243 doi:10.1029/2002JB002287.
- 244 Higgs, K.T., Boyd, J.D., and Williams, B.P.J., 2014, An Early Devonian age for the Bulls
245 Head Formation: Lower Old Red Sandstone: Dingle Peninsula: Irish Journal of Earth
246 Science, v. 32, p. 55–70, doi:10.3318/IJES.2014.32.55.

- 247 Horne, R., 1974, The lithostratigraphy of the late Silurian to early Carboniferous of the
248 Dingle Peninsula, County Kerry: Geological Survey of Ireland Bulletin, v. 1, p. 395–
249 428.
- 250 Jeffery, G.B., 1922, The motion of ellipsoidal particles immersed in a viscous fluid:
251 Proceedings of the Royal Society of London. Series A, Containing Papers of a
252 Mathematical and Physical Character, v. 102, p. 161–179,
253 doi:10.1098/rspa.1922.0078.
- 254 Maxwell, J., 1962, Origin of slaty and fracture cleavage in the Delaware Water Gap area,
255 New Jersey and Pennsylvania, *in* Engel, A.E.J., James, H.L., and Leonard, B.F.,
256 eds., Geological Society of America Petrologic Studies: A volume to honor A.F.
257 Buddington: Boulder, Colorado, Geological Society of America, p. 281–311,
258 doi:10.1130/Petrologic.1962.281.
- 259 Meere, P.A., and Mulchrone, K.F., 2003, The effect of sample size on geological strain
260 estimation from passively deformed clastic sedimentary rocks: Journal of Structural
261 Geology, v. 25, p. 1587–1595, doi:10.1016/S0191-8141(03)00007-5.
- 262 Meere, P.A., and Mulchrone, K.F., 2006, Timing of deformation within Old Red
263 Sandstone lithologies from the Dingle Peninsula, SW Ireland: Journal of the
264 Geological Society, v. 163, p. 461–469, doi:10.1144/0016-764905-099.
- 265 Meere, P.A., Mulchrone, K.F., Sears, J.W., and Bradway, M.D., 2008, The effect of non-
266 passive clast behaviour in the estimation of finite strain in sedimentary rocks: Journal
267 of Structural Geology, v. 30, p. 1264–1271, doi:10.1016/j.jsg.2008.06.008.

- 268 Meere, P.A., Mulchrone, K.F., and Timmerman, M., 2013, Shear folding in low-grade
269 metasedimentary rocks: Reverse shear along cleavage at a high angle to the
270 maximum compressive stress: *Geology*, v. 41, p. 879–882, doi:10.1130/G34150.1.
- 271 Mulchrone, K.F., 2007a, Shape fabrics in populations of rigid objects in 2D: Estimating
272 finite strain and vorticity: *Journal of Structural Geology*, v. 29, p. 1558–1570,
273 doi:10.1016/j.jsg.2007.06.006.
- 274 Mulchrone, K.F., 2007b, An analytical solution in 2D for the motion of rigid elliptical
275 particles with a slipping interface under a general deformation: *Journal of Structural*
276 *Geology*, v. 29, p. 950–960, doi:10.1016/j.jsg.2007.03.008.
- 277 Mulchrone, K.F., and Meere, P.A., 2015, Shape fabric development in rigid clast
278 populations under pure shear: The influence of no-slip versus slip boundary
279 conditions: *Tectonophysics*, v. 659, p. 63–69, doi:10.1016/j.tecto.2015.08.003.
- 280 Mulchrone, K.F., O’Sullivan, F., and Meere, P.A., 2003, Finite strain estimation using the
281 mean radial length of elliptical objects with bootstrap confidence intervals: *Journal*
282 *of Structural Geology*, v. 25, p. 529–539, doi:10.1016/S0191-8141(02)00049-4.
- 283 Mulchrone, K.F., McCarthy, D.J., and Meere, P.A., 2013, Mathematica code for image
284 analysis, semi-automatic parameter extraction and strain analysis: *Computers &*
285 *Geosciences*, v. 61, p. 64–70, doi:10.1016/j.cageo.2013.08.001.
- 286 Paterson, S.R., and Tobisch, O.T., 1993, Pre-lithification structures, deformation
287 mechanisms, and fabric ellipsoids in slumped turbidites from the Pigeon Point
288 Formation, California: *Tectonophysics*, v. 222, p. 135–149, doi:10.1016/0040-
289 1951(93)90045-L.

- 290 Phillips, C.A., and Alsop, G.I., 2000, Post-tectonic clastic dykes in the Dalradian of
291 Scotland and Ireland: implications for delayed lithification and deformation of
292 sediments: *Geological Journal*, v. 35, p. 99–110, doi:10.1002/1099-
293 1034(200004/06)35:2<99::AID-GJ844>3.0.CO;2-X.
- 294 Powell, C.McA., 1979, A morphological classification of rock cleavage: *Tectonophysics*,
295 v. 58, p. 21–34, doi:10.1016/0040-1951(79)90320-2.
- 296 Sorby, H.C., 1849, On the Origin of Slaty Cleavage: *Proceedings of the Yorkshire*
297 *Geological Society*, v. 3, p. 300–312, doi:10.1144/pygs.3.300.
- 298 Todd, S.P., 2000, Taking the roof off a suture zone: basin setting and provenance of
299 conglomerates in the ORS Dingle Basin of SW Ireland: *Geological Society of*
300 *London Special Publications*, v. 180, p. 185–222,
301 doi:10.1144/GSL.SP.2000.180.01.10.
- 302 Van den Driessche, J., and Brun, J.-P., 1987, Rolling structures at large shear strain:
303 *Journal of Structural Geology*, v. 9, p. 691–704, doi:10.1016/0191-8141(87)90153-2.
- 304 Vernon, R.H., 1998, Chemical and volume changes during deformation and prograde
305 metamorphism of sediments, *in* Treloar, P.J., and O'Brien, P.J., eds., *What drives*
306 *metamorphism and metamorphic reactions?: Geological Society of London Special*
307 *Publications*, 138, p. 215–246, doi:10.1144/GSL.SP.1996.138.01.13.
- 308 Vernon, R.H., 2004, *A practical guide to rock microstructures*: Cambridge, UK,
309 Cambridge University Press, 594pp, doi:10.1017/CBO9780511807206.
- 310 Waldron, J.W.F., and Gagnon, J.-F., 2011, Recognizing soft-sediment structures in
311 deformed rocks of orogens: *Journal of Structural Geology*, v. 33, p. 271–279,
312 doi:10.1016/j.jsg.2010.06.015.

Wood, D.S., 1974, Current views on the development of slaty cleavage: Annual Reviews
of Earth Science, v. 2, p. 369–401, doi:10.1146/annurev.ea.02.050174.002101.

FIGURE CAPTIONS

Figure 1. Geological map of the northwestern Dingle Peninsula (Ireland) with an equal
area projection of structural data for the Ballyferriter Syncline in the Wine Strand area
demonstrating anticlockwise transection of the calculated fold axis (x) by the associated
tectonic fabric (S_1). Filled points are poles to bedding, solid great circles are S_1 planes.
Gp.—Group; Fm.—Formation.

Figure 2. Meso-structural field evidence of the contrasting competencies between
competent extra-formational and incompetent intra-formational (rip-up) clasts, and the
surrounding incompetent sand grade matrix. A: Field image of deformed conglomerate
with competent jasper (j), mudstone (m), and volcanic clasts (v) in addition to ‘rip-up’
incompetent red mudstone clasts (r-u) set in a sand grade matrix. Note wrapping of
cleavage fabric (S_1) around jasper clast while the sand matrix is seen to ‘bulge’ into the
less competent ‘rip-up’ clast (23-mm-diameter coin for scale). B: View of more
competent volcanic clast projecting into less competent ‘rip-up’ clast, note localized
development of dissolution seams (ds) associated with high tectonic stress concentrations
at the apices of the more competent volcanic clast. Also note the highly angular nature of
the sandstone matrix clasts, the absence of cleavage domains and a clast shape fabric
parallel to S_1 in the lower third of the image. C: Bulging (arrows) of coarse-grained
sandstone and pebble conglomerate matrix into mudstone rip-up clast. D: Strong
alignment of ‘rip-up’ clasts parallel to the cleavage fabric and at a high angle to the

bedding fabric (S_0). E: View of mullioned contact across the cleavage (S_1), detail shows reduction spot in approximately the xz plane of the finite strain ellipsoid with an R_s value of ~3.5. F: View of mudstone/conglomerate mullion contact in the plane of cleavage, note lobate nature of contact along the mullion lineation.

Figure 3. Photomicrographs and electron microprobe Si concentration maps of siltstone (Siltst.) close to a siltstone/sandstone (Sst.) boundary (A), note lack of silica depleted dissolution seams in the siltstone (sample 24–6–13–3), and siltstone close to a siltstone/sandstone boundary with a very large volcanic clast impinging on the siltstone (B) resulting in the very localized development of dissolution seams (DS) now outlined by Mn-oxides (sample 24–6–13–1b). C—chlorite, M—muscovite, P—plagioclase, Q—quartz, V—volcanic clast.

Figure 4. A: Plot of finite strain (R_s) estimates with 95% confidence interval error bars determined using mean radial length analysis of sedimentary clasts versus deviation of principle strain axis ϕ from cleavage (S_1). B: Plot of variation in clast fabric intensity versus bulk strain (R_s) for slipping and sticking clast/matrix behaviors.







